

Ramon Subirós- Funosas<sup>a</sup>, Ayman El-Faham<sup>a,b</sup>, Fernando Albericio<sup>a,c</sup><sup>a</sup> Institute for Research in Biomedicine, Barcelona Science Park, Baldiri Reixac 10, 08028-Barcelona, Spain<sup>b</sup> Department of Chemistry, Faculty of Science, Alexandria University, Ibrahimia 21321, Alexandria, Egypt<sup>c</sup> Department of Organic Chemistry, University of Barcelona, Martí i Franquès 1-11, 08028-Barcelona, Spain
[ramon.subiros@irbbarcelona.org](mailto:ramon.subiros@irbbarcelona.org), [aymanel\\_faham@hotmail.com](mailto:aymanel_faham@hotmail.com), [albericio@pcb.ub.cat](mailto:albericio@pcb.ub.cat)

## 1. Introduction

Benzotriazole-based additives, especially HOBT<sup>1</sup> and HOAt<sup>2</sup>, have been widely used along with carbodiimides or stand-alone coupling reagents, in order to suppress racemization or other side reactions. Nonetheless, recent studies<sup>3</sup> revealed the explosive properties of such compounds that led to their classification as Class 1 (transport of explosive substances), thereby increasing transport restrictions. The need for a new family of safe and potent additives has therefore become of marked importance in peptide synthesis. In that scenario, *ethyl 2-hydroxyimino-2-cyanoacetate* (Oxyma pure, Fig.1), which had been previously reported<sup>4</sup>, has been evaluated in terms of racemization control, efficiency in difficult couplings and safety profile.




Figure 1. Structure of oxyma pure

## 3. Efficiency in difficult couplings

The performance of Oxyma Pure in obtaining a modified Leu-enkephalin pentapeptide (H-Tyr-Aib-Aib-Phe-Leu-NH<sub>2</sub>)<sup>6</sup> was tested. This model peptide, containing Aib ( $\alpha$ -aminoisobutyric acid) residues, proved to be the most demanding one, and so bigger differences among HOAt, HOBT and Oxyma Pure arose in terms of ratio pentapeptide/deletion peptides observed. Incorporation of Tyr and Aib to a previously elongated H-Aib-Phe-Leu-resin tripeptide was followed, using DIC/additive/Fmoc-amino acid (3eq) with a 3 min PT to ensure formation of the corresponding active esters. Different coupling conditions were studied (Table 3). In all cases HOBT gave rise to the lowest percentages of pentapeptide obtained (des-Tyr and tripeptide were detected even after 1 hour coupling). Oxyma Pure showed greater efficiency than HOAt in every experiment.

Table 3. H-Tyr-Aib-Aib-Phe-Leu-NH<sub>2</sub>

Coupling time	additive	Pentapeptide (%)	des-Aib (%)	des-Tyr (%)	Tripeptide (%)
30 minutes	HOAt	11.3	86.1	1.8	0.8
	HOBT	3.0	91.0	0.9	5.1
	Oxyma Pure	28.0	70.5	1.1	0.4
1 hour	HOAt	28.7	71.3	1.7	1.7
	HOBT	9.8	86.9	1.6	1.7
	Oxyma Pure	55.7	44.3	--	--
1 hour double coupling	HOAt	55.0	45.0	--	--
	HOBT	18.9	80.6	0.5	--
	Oxyma Pure	69.0	31.0	--	--

## 4. Differential Scanning Calorimetry Assays

For a compound to be explosive, a fast release of pressure is necessary when decomposing. To address this issue, Differential Scanning Calorimetry (DSC) tests comparing HOAt, HOBT and Oxyma Pure were carried out. In this experiment, information is given concerning the kinetics of decomposition. The sample is heated from 30°C to 300°C at a constant heating rate of 10°C/min in a closed crucible with N<sub>2</sub> flow. A diagram with released dH vs temperature/ time is obtained. Benzotriazole-based additives displayed similar profiles, (Fig.2 and Fig.3) revealing that decomposition takes place in a short period of time, whereas Oxyma Pure (Fig.4) decomposed much slower.

Fig 2. HOAt




Fig 3. HOBT





Fig 4. Oxyma Pure



## 5. Adiabatic Calorimetry Assays

Complementary to the above mentioned DSC experiments are the adiabatic calorimetry assays run on an Accelerating Rate Calorimeter (ARC) applying the "heat-wait-see" method. The sensitivity threshold was set to 0.02°C/min. Data is given regarding the pressure released in the decomposition process (temperature in red or pressure in blue vs time). Whereas in the experiment with the benzotriazole-based additives HOAt and HOBT (Fig.5 and Fig.6) 170-180 bar pressure is detected, due to N<sub>2</sub> release, in the case of Oxyma Pure only 60 bar pressure (Fig.7) was observed.

Fig 5. HOAt




Fig 6. HOBT





Fig 7. Oxyma Pure



## Conclusions

Results on Oxyma Pure have shown that:

- The suppression of racemization is at least comparable to HOAt
- The obtention of sterically hindered peptides is more efficient than HOAt and HOBT.
- The likelihood of an explosion can not be compared to HOAt or HOBT, because it decomposes slowly releasing low pressure

## References

- 1 König, W.; Geiger, R.; *Chem.Ber.* **1970**, *103*, 788-798 and 2034-2040
- 2 Carpino, L.A.; *J.Am.Chem.Soc.* **1993**, *115*, 4397-4398
- 3 Wehrstedt et al., *J. Hazard. Mat.* **2005**, *A126*, 1-7
- 4 a) Izdebski, J., *Pol.J.Chem.* **1979**, *53*, 1049-1057; b) Itoh, M., *Bull.Chem.Soc. Japan*, **1973**, *46*, 2219-2221
- 5 a) El-Faham, A.; Albericio, F. *Org. Lett.* **2007**, *9*, 4475-4477.; b) El-Faham, A.; Albericio, F. *J. Org. Chem.*, **2008**, *73*, 2731-2737.
- 6 Carpino, L.A.; El-Faham, A.; Minor, C.A.; Albericio, F. *J. Chem. Soc., Chem.Comm.* **1994**, pp. 201-203

## Acknowledgements

This work was partially supported by CICYT (CTQ2006-03794/BQU), the Generalitat de Catalunya (2005SGR 00662), the Institute for Research in Biomedicine, and the Barcelona Science Park. RS-F thanks Ministerio de Educación y Ciencia for a FPU PhD fellowship. We also thank the Calorimetry Platform at Barcelona Science Park for their support in the DSC and ARC experiments.

2 Bergman St. Rehovot 76705, ISRAEL

Tel: +972 8 9466040 Fax: +972 8 9466402 [www.peptide-and-dna.com](http://www.peptide-and-dna.com)

marina@luxembourg.co.il